

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 April 2004 (15.04.2004)

PCT

(10) International Publication Number
WO 2004/031182 A1

(51) International Patent Classification⁷: C07D 409/14,
403/14, 403/12, 471/04, 471/04, 221/00, 209/00

(74) Agent: SERONO INTERNATIONAL S.A. INTEL-
LECTUAL PROPERTY; 12, Chemin des Aulx, CH-1228
Plan-les-Ouates (CH).

(21) International Application Number:
PCT/EP2003/050640

(22) International Filing Date:
19 September 2003 (19.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/412,308 20 September 2002 (20.09.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): AP-
PLIED RESEARCH SYSTEMS ARS HOLDING N.V.
[NL/NL]; Pietermaai 15, Curacao (AN).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): MAGAR, Sharad
[US/US]; 20 Harrison Road, Canton, MA 02021 (US).
GOUTOPOULOS, Andreas [GR/US]; 250 New-
bury Street, Boston, MA 02116 (US). LIAO, Yihua
[CN/US]; 40 Highview Street, Westwood, MA 02090
(US). SCHWARZ, Matthias [CH/CH]; Avenue Tronchet
32B, CH-1226 Thônex (CH). RUSSELL, Thomas, J.
[GB/GB]; Cummor Road 149, Boars Hill, Oxford OX1
5JS (GB).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 2004/031182 A1

(54) Title: PIPERAZINE DERIVATIVES AND METHODS OF USE

(57) Abstract: The invention provides 2-carboxamide piperazine compounds of formula I, R³. I N¹ N² ~ R²(O)Z- o R⁴I wherein R¹, R², R³ and R⁴ are as defined in the claims and methods of treatment and pharmaceutical compositions that utilize or comprise one or more such compounds. Compounds of the invention are useful for the treatment of mammalian infertility.

PIPERAZINE DERIVATIVES AND METHODS OF USE

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to 2-carboxamide piperazine compounds, and methods of treatment and pharmaceutical compositions that utilize or comprise one or more such compounds. Compounds of the invention are useful for the treatment of mammalian infertility.

2. Background

Luteinizing Hormone (LH) and Follicle Stimulating Hormone (FSH) are produced by the anterior pituitary gland and are involved in mammalian reproductive processes. The glycoprotein family of pituitary hormones such as FSH, LH and the thyrotropic hormone (TSH) relatively large heterodimeric proteins that contain carbohydrate moieties. They have a common α -subunit and distinct β -subunits (1-4) providing receptor recognition and specificity.

LH is released from the anterior pituitary gland under the influence of gonadotropin releasing hormone and progesterins. In the female LH stimulates ovulation and is the major hormone involved in the regulation of progesterone secretion by the corpus luteum. In the male, LH stimulates Leydig cells to secrete androgens, particularly testosterone.

FSH is released from the anterior pituitary under the influence of the gonadotrophin releasing hormone, and oestrogens and from the placenta during pregnancy. FSH acts on the ovaries stimulating maturation of follicles and regulates secretion of oestrogens. In the male, FSH is responsible for the integrity of the seminiferous tubules and acts on Sertoli cells to support gametogenesis.

The LH and FSH receptors belong to the superfamily of G-protein coupled receptors, which are complex transmembrane proteins characterized by seven

hydrophobic helices. The LH and FSH receptors also share close sequence homology (approximately 40%). The receptors' extracellular domains bind to their respective hormones with high affinity and specificity. The intracellular portions of FSH and LH receptors are coupled to a Gs protein. Receptor activation upon the hormonal interaction with the extracellular domain results in a cascade of events that leads to specific biological effects.

LH and FSH have been used for the treatment of female infertility and spermatogenesis disorders. See U.S. Patents 5,767,067; 5,639,639; and 5,017,557.

However, those therapies have some notable shortcomings. For instance, current FSH treatment is limited by lack of oral bioavailability, high costs and need of close medical personnel supervision throughout an administration protocol.

Certain non-peptidic FSH agonists also have been disclosed. See, for instance, U.S. Patent 6,235,755; 6,423,723; WO 00/08015; and WO 02/09706.


Thus, it would be desirable to have new agents and methods to treat infertility in mammals.

SUMMARY OF THE INVENTION

We have now found that 2-carboxamide piperazine compounds are potent Follicle Stimulating Hormone receptor (FSH) agonists. Compounds of the invention are particularly useful for treatment of infertility in mammals.

More specifically, in a preferred aspect of the invention, preferred 2-carboxamide piperazine compounds of the invention include those of the following Formula I:

3



I


wherein R¹ and R² are independently selected from the group comprising or consisting of hydrogen, C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₂-C₁₂-alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group, an acyl moiety, C₁-C₁₂-alkyl aryl, C₁-C₁₂-alkyl heteroaryl, C₂-C₁₂-alkenyl aryl, C₂-C₁₂-alkenyl heteroaryl, C₂-C₁₂-alkynyl aryl, C₂-C₁₂-alkynyl heteroaryl, C₁-C₁₂-alkyl cycloalkyl, C₁-C₁₂-alkyl heterocycloalkyl, C₂-C₁₂-alkenyl cycloalkyl, C₂-C₁₂-alkenyl heterocycloalkyl, C₂-C₁₂-alkynyl cycloalkyl, C₂-C₁₂-alkynyl heterocycloalkyl, alkoxycarbonyl, aminocarbonyl, C₁-C₁₂-alkyl carboxy, C₁-C₁₂-alkyl acyl, aryl acyl, heteroaryl acyl, C₃-C₈-(hetero)cycloalkyl acyl, C₁-C₁₂-alkyl acyloxy, C₁-C₁₂-alkyl alkoxy, C₁-C₁₂-alkyl alkoxycarbonyl, C₁-C₁₂-alkyl aminocarbonyl, C₁-C₁₂-alkyl acylamino, acylamino, C₁-C₁₂-alkyl ureido, C₁-C₁₂-alkyl carbamate, C₁-C₁₂-alkyl amino, C₁-C₁₂-alkyl ammonium, C₁-C₁₂-alkyl sulfonyloxy, C₁-C₁₂-alkyl sulfonyl, C₁-C₁₂-alkyl sulfinyl, C₁-C₁₂-alkyl sulfanyl, C₁-C₁₂-alkyl sulfonylamino, or C₁-C₁₂-alkyl aminosulfonyl, and preferably with at least one of R¹ and R² being other than hydrogen;

R³ is C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₂-C₁₆-alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group, an acyl moiety, C₁-C₁₆-alkyl aryl, C₁-C₁₆-alkyl heteroaryl, C₂-C₁₆-alkenyl aryl, C₂-C₁₆-alkenyl heteroaryl, C₂-C₁₆-alkynyl aryl, C₂-C₁₆-alkynyl heteroaryl, C₁-C₁₆-alkyl cycloalkyl, C₁-C₁₆-alkyl heterocycloalkyl, C₂-C₁₆-alkenyl cycloalkyl, C₂-C₁₆-alkenyl heterocycloalkyl, C₂-C₁₆-alkynyl cycloalkyl, C₂-C₁₆-alkynyl heterocycloalkyl, alkoxycarbonyl, aminocarbonyl, C₁-C₁₆-alkyl carboxy, C₁-

C₁₆-alkyl acyl, aryl acyl, heteroaryl acyl, C₃-C₈-(hetero)cycloalkyl acyl, C₁-C₁₆-alkyl acyloxy, C₁-C₁₆-alkyl alkoxy, C₁-C₁₆-alkyl alkoxy carbonyl, C₁-C₁₆-alkyl aminocarbonyl, C₁-C₁₆-alkyl acylamino, acylamino, C₁-C₁₆-alkyl ureido, C₁-C₁₆-alkyl carbamate, C₁-C₁₆-alkyl amino, C₁-C₁₆-alkyl ammonium, C₁-C₁₆-alkyl sulfonyloxy, C₁-C₁₆-alkyl sulfonyl, C₁-C₁₆-alkyl sulfinyl, C₁-C₁₆-alkyl sulfanyl, C₁-C₁₆-alkyl sulfonylamino, or C₁-C₁₆-alkyl aminosulfonyl;

R⁴ is C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₂-C₁₂-alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group or amino group; and pharmaceutically acceptable salts thereof.

Generally preferred compounds of the invention include those where the substituent R⁴ of Formula I is heteroaryl, particularly a sulfur-ring substituent especially such as compounds of the following Formula II:



II


wherein R¹, R² and R³ are the same as defined above in Formula I;

each R⁵ is independently halogen, hydroxy or the same as defined for R¹;

m is an integer of from 0 (where available thienyl ring positions are hydrogen-substituted) to 4, and preferably m is 0, 1, 2 or 3, more preferably 0, 1 or 2; and pharmaceutically acceptable salts thereof.

Compounds of Formula II having a 2-thienyl moiety are typically preferred, i.e.

compounds of the following Formula III:



III

wherein R^1 , R^2 , R^3 , R^5 and m are the same as defined for Formula II above; and pharmaceutically acceptable salts thereof.

As mentioned above, in each of Formulae I through III, preferably at least one of R^1 and R^2 is other than hydrogen, and more preferably, one of R^1 and R^2 is hydrogen and the other a non-hydrogen substituent. Preferred non-hydrogen R^1 and R^2 substituents include optionally substituted aryl and optionally substituted heteroaryl groups such as optionally substituted phenyl, naphthyl, pyridyl, and fused aryl and heteroaryl ring systems such as benzoimidazole, pyridyl-substituted benzoimidazole, carbazole and the like. Especially preferred non-hydrogen R^1 and R^2 substituents include those that comprise an optionally substituted carbazolyl, an optionally substituted tetrahydro-beta-carbolinyl or an optionally substituted benzimidazolyl moiety, such as a 1-benzimidazolyl which may be N-substituted by C_1 - C_3 alkyl, 2,3,4,9-tetrahydro-beta-carbolin-6yl-1-one and 9-ethyl-9H-carbazol-3yl or the like.

In Formulae I through III, preferred R^3 groups contain 4 or more atoms, particularly in a sequential chain, even more preferably 5, 6, 7, 8, 9, or 10 atoms in an extended chain such as an alkyl chain or ester, amide or other moiety. Alkyl groups, including n-alkyl groups, that contain 4, 5, 6, 7, 8, 9 or 10 carbon atoms are particularly preferred R^3 substituents.

A preferred embodiment of the invention, is a piperazine derivative according to Formula I wherein R^1 is H; R^2 is selected from optionally substituted aryl, optionally substituted heteroaryl, optionally substituted saturated or unsaturated 3-8-membered cycloalkyl and optionally substituted heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group; R^3 is selected from optionally substituted C_1 - C_{16} -alkyl, optionally substituted C_2 - C_{16} -alkenyl, optionally substituted C_2 - C_{16} -alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N and O, optionally substituted monocyclic aryl, optionally substituted monocyclic heteroaryl, optionally substituted saturated or optionally substituted unsaturated 3-8-membered monocyclic cycloalkyl, optionally substituted monocyclic heterocycloalkyl, an acyl moiety, optionally substituted C_1 - C_{16} -alkyl aryl, optionally substituted C_1 - C_{16} -alkyl heteroaryl, optionally substituted C_2 - C_{16} -alkenyl aryl, optionally substituted C_2 - C_{16} -alkenyl heteroaryl, optionally substituted C_2 - C_{16} -alkynyl aryl, optionally substituted C_2 - C_{16} -alkynyl heteroaryl, optionally substituted C_1 - C_{16} -alkyl cycloalkyl, optionally substituted C_1 - C_{16} -alkyl heterocycloalkyl, optionally substituted C_2 - C_{16} -alkenyl cycloalkyl, optionally substituted C_2 - C_{16} -alkenyl heterocycloalkyl, optionally substituted C_2 - C_{16} -alkynyl cycloalkyl, optionally substituted C_2 - C_{16} -alkynyl heterocycloalkyl, optionally substituted alkoxycarbonyl, optionally substituted aminocarbonyl, optionally substituted C_1 - C_{16} -alkyl carboxy, optionally substituted C_1 - C_{16} -alkyl acyl, optionally substituted aryl acyl, optionally substituted heteroaryl acyl, optionally substituted C_3 - C_8 -(hetero)cycloalkyl acyl, optionally substituted C_1 - C_{16} -alkyl acyloxy, optionally substituted C_1 - C_{16} -alkyl alkoxy, optionally substituted C_1 - C_{16} -alkyl alkoxycarbonyl, optionally substituted C_1 - C_{16} -alkyl aminocarbonyl, optionally substituted C_1 - C_{16} -alkyl acylamino, optionally substituted acylamino, optionally substituted C_1 - C_{16} -alkyl sulfinyl, optionally substituted C_1 - C_{16} -alkyl sulfanyl, optionally substituted C_1 - C_{16} -alkyl ureido, optionally substituted C_1 - C_{16} -alkyl carbamate, optionally substituted C_1 - C_{16} -alkyl amino and optionally substituted C_1 - C_{16} -alkyl ammonium; R^4 is selected from optionally substituted C_1 - C_{12} -alkyl, optionally substituted C_2 - C_{12} -alkenyl, optionally substituted C_2 - C_{12} -alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O and S, optionally substituted aryl, optionally substituted heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein

said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group, and optionally substituted amino group.

A particularly preferred embodiment, is a piperazine derivative according to Formulae I to III wherein R^1 is H and R^2 is selected from optionally substituted aryl and optionally substituted heteroaryl, wherein optionally substituted aryl and optionally substituted heteroaryl can be optionally polycyclic and include optionally substituted phenyl, optionally substituted benzimidazolyl, optionally substituted carbazolyl and optionally substituted tetrahydro carbolinyl.

A further preferred embodiment, is a piperazine derivative according to Formulae I to III, wherein R^1 is H and R^2 is selected from optionally substituted aryl and optionally substituted heteroaryl, wherein optionally substituted aryl and optionally substituted heteroaryl are bi-cyclic or tricyclic and include optionally substituted benzimidazolyl, optionally substituted carbazolyl and optionally substituted tetrahydro carbolinyl.

A particularly preferred embodiment, is a piperazine derivative according to Formulae I to III, wherein R^1 is H and R^3 is selected from optionally substituted C_1 - C_8 -alkyl, including pentyl, hexyl, heptyl and octyl; optionally substituted C_1 - C_8 -alkyl acyloxy, including acetic acid ethyl ester; optionally substituted C_1 - C_8 -alkyl sulfanyl, including methylsulfanyl propyl; optionally substituted C_1 - C_8 -alkyl heteroaryl, including imidazolyl propyl and ethyl furanylmethyl; optionally substituted aminocarbonyl, including ethoxy ethyl formamide, formylamino-acetic acid ethyl ester and imidazolyl propylamide; optionally substituted C_1 - C_8 -alkyl aminocarbonyl, including ethylamide and pentylamide and optionally substituted carbonyl, including pentanoyl.

A further preferred embodiment, is a piperazine derivative according to Formulae I to III, wherein R^1 is H and R^3 is selected from optionally substituted C_1 - C_8 -alkyl, including pentyl, hexyl, heptyl and octyl and optionally substituted C_1 - C_8 -alkyl aminocarbonyl, including ethylamide and pentylamide.

A particularly preferred embodiment, is a piperazine derivative according to Formula I wherein R^1 is H and R^4 is selected from optionally substituted C_1 - C_6 -alkyl, wherein said alkyl chain may be interrupted by a nitrogen atom, including butane; optionally substituted amino group, including dimethyl amine; optionally substituted aryl, including optionally substituted phenyl such as phenyl and fluorophenyl; optionally substituted heteroaryl, including thienyl and methyl imidazole.

A further preferred embodiment, is a piperazine derivative according to Formula I wherein R^1 is H and R^4 is selected from optionally substituted aryl, including optionally substituted phenyl such as fluorophenyl; optionally substituted heteroaryl, including thiophene and methyl imidazole.

Generally preferred R^4 substituents of Formula I are optionally substituted alkyl, optionally substituted aryl and optionally substituted heteroaryl groups, such as optionally substituted C_{1-12} alkyl, optionally substituted aryl such as optionally substituted phenyl e.g. phenyl having one or more ring substituents of halogen, nitro, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, and the like, and optionally substituted heteroaryl, particularly thiophene and furane.

The invention also includes compounds and use of optically active compounds of the above Formulae I through III, particularly compounds of the above Formulae I through III where a single stereoisomer of a chiral compound is present in an enantiomeric excess, e.g. where a single stereoisomer is present in an amount of at least 70 mole percent relative to other stereoisomer(s), more preferably where one stereoisomer is present in an amount of at least about 80, 85, 90, 92, 93, 94, 95, 96, 97, 98 or 99 mole percent relative to other stereoisomer(s). Particularly preferred is where a stereoisomer of the depicted chiral carbon (i.e. the chiral piperazine ring carbon indicated by * in each Formulae I, II and III) is present in an enantiomeric excess, such as the amounts specified above relative to other stereoisomer(s).

Carboxamide piperazine compounds of the invention are useful for treatment of infertility in male and female mammals, particularly humans. Therapeutic methods of the invention in general comprise administering an effective amount of one or more 2-carboxamide piperazine compounds as disclosed herein to a mammal in need thereof, such as a mammal suspected of suffering from infertility, particularly a human suspected of suffering from infertility. Compounds of the invention will be useful for treatment of infertility conditions currently treated with FSH and/or LH, including female infertility and male spermatogenesis disorders.

Additionally, in contrast to current protein therapeutics such as FSH, compounds of the invention can be administered orally and without extensive medical specialist supervision.

Preferred compounds of the invention exhibit good agonist activity in standard Follicle Stimulating Hormone (FSH) assays, such as the assay of Example 2 which follows.

It also has been found that 2-carboxamide piperazine compounds of the invention exhibit good inhibition activity against phosphodiesterase PDE4, adenosine transporters, and prostanoid receptors. Accordingly, methods are provided for treatment of diseases and disorders associated with phosphodiesterase PDE4, adenosine transporters, and prostanoid receptors, which methods in general comprise administration of an effective amount of one or more 2-carboxamide piperazine compounds to a patient (e.g. mammal, such as human or other primate) in need of such treatment.

In a further aspect, the invention provides a use of a carboxamide piperazine compound as disclosed herein, particularly a compound of any one of Formulae I through III for the treatment or prevention (including prophylactic treatment) of a disease or condition as disclosed herein, including treatment of infertility in male and female mammals, particularly humans, such as female infertility and male spermatogenesis disorders, as well as treatment of diseases and disorders associated with phosphodiesterase PDE4, adenosine transporters, and prostanoid receptors.

In a yet further aspect, the invention provides use of a carboxamide piperazine compound as disclosed herein, particularly a compound of any one of Formulae I through III, for the preparation of a medicament for the treatment or prevention (including prophylactic treatment) of a disease or condition as disclosed herein, including treatment of infertility in male and female mammals, particularly humans, such as female infertility and male spermatogenesis disorders, as well as treatment of diseases and disorders associated with phosphodiesterase PDE4, adenosine transporters, and prostanoid receptors.

Preferred methods of the invention including identifying and/or selecting a subject (e.g. mammal, particularly human) that is susceptible or suffering from a condition disclosed herein, and thereafter administering to the identified and selected subject one or more carboxamide piperazine compounds of the invention, particularly a subject that is identified and selected as being susceptible to or suffering from infertility, such as female infertility and male spermatogenesis disorders in humans, as well as subjects susceptible to or suffering from a disease or disorder associated with phosphodiesterase PDE4, adenosine transporters, and prostanoid receptors.

The invention also provides pharmaceutical compositions that comprise one or more 2-carboxamide piperazine compounds of the invention and a suitable carrier for the compositions. Other aspects of the invention are disclosed *infra*.

DETAILED DESCRIPTION OF THE INVENTION

We have now discovered that substituted carboxamide piperazine compounds, including compounds of the above Formulae I, II and III, that are useful for treatment of infertility in mammals, including female and male humans.

The following paragraphs provide definitions of the various chemical moieties that make up the compounds according to the invention and are intended to apply uniformly through-out the specification and claims, unless an otherwise expressly set out definition provides a broader definition.

"C₁-C₁₂-alkyl" refers to monovalent alkyl groups having 1 to 12 carbon atoms. This term is exemplified by groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, *tert*-butyl, n-hexyl and the like. By analogy, "C₁-C₁₆-alkyl" refers to monovalent alkyl groups having 1 to 16 carbon atoms, "C₁-C₈-alkyl" refers to monovalent alkyl groups having 1 to 8 carbon atoms and "C₁-C₆-alkyl" refers to monovalent alkyl groups having 1 to 6 carbon atoms.

"Aryl" refers to an unsaturated aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring or "monocyclic ring" (e.g., phenyl), multiple condensed rings or "polycyclic ring", including for example "bi-cyclic" ring (e.g., naphthyl), "tri-cyclic" ring (e.g. 9-carbazolyl) or other linked multiple rings (e.g., bi-phenyl). Preferred aryl include phenyl, naphthyl, anthracenyl, acenaphthyl, phenanthrenyl and the like. The expression "fused aryl" means an aryl that is fused with further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group(s), preferably 1 or 2.

"C₁-C₁₂-alkyl aryl" refers to C₁-C₁₂-alkyl groups having an aryl substituent, including benzyl, phenethyl and the like. By analogy, "C₁-C₁₆-alkyl aryl" refers to C₁-C₁₆-alkyl groups having an aryl substituent and "C₁-C₈-alkyl aryl" refers to C₁-C₈-alkyl groups having an aryl substituent.

"Heteroaryl" refers to a monocyclic heteroaromatic, or a bicyclic or a tricyclic fused-ring heteroaromatic group. Particular examples of heteroaromatic groups include optionally substituted pyridyl, pyrrolyl, furyl, thienyl, imidazolyl, benzoxazole, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,3,4-triazinyl, 1,2,3-triazinyl, benzofuryl, [2,3-dihydro]benzofuryl, isobenzofuryl, benzothieryl, benzotriazolyl, isobenzothieryl, indolyl, isoindolyl, 3H-indolyl, benzimidazolyl, imidazo[1,2-a]pyridyl, benzothiazolyl, benzoxa-zolyl, quinoliziny, quinazolinyl, pthalaziny, quinoxaliny, cinnolinyl, naphthyridiny, pyrido[3,4-b]pyridyl, pyrido[3,2-b]pyridyl, pyrido[4,3-b]pyridyl, quinolyl, isoquinolyl, tetrazolyl, 5,6,7,8-

tetrahydroquinolyl, 5,6,7,8-tetrahydroisoquinolyl, purinyl, pteridinyl, carbazolyl, xanthenyl or benzoquinolyl.

"C₁-C₁₂-alkyl heteroaryl" refers to C₁-C₁₂-alkyl groups having a heteroaryl substituent, including 2-furylmethyl, 2-thienylmethyl, 2-(1H-indol-3-yl)ethyl and the like. By analogy, "C₁-C₁₆-alkyl heteroaryl" refers to C₁-C₁₆-alkyl groups having a heteroaryl substituent and "C₁-C₈-alkyl heteroaryl" refers to C₁-C₈-alkyl groups having an heteroaryl substituent.

"C₂-C₁₂-alkenyl" refers to alkenyl groups preferably having from 2 to 12 carbon atoms and having at least 1 or 2 sites of alkenyl unsaturation. By analogy, "C₂-C₁₆-alkenyl" refers to alkenyl groups preferably having from 2 to 16 carbon atoms and having at least 1 or 2 sites of alkenyl unsaturation. Preferable alkenyl groups include ethenyl (-CH=CH₂), n-2-propenyl (allyl, -CH₂CH=CH₂) and the like.

"C₂-C₁₂-alkenyl aryl" refers to C₂-C₁₂-alkenyl groups having an aryl substituent, including 2-phenylvinyl and the like. By analogy, "C₂-C₁₆-alkenyl aryl" refers to C₂-C₁₆-alkenyl groups having an aryl substituent

"C₂-C₁₂-alkenyl heteroaryl" refers to C₂-C₁₂-alkenyl groups having a heteroaryl substituent, including 2-(3-pyridinyl)vinyl and the like. By analogy, "C₂-C₁₆-alkenyl heteroaryl" refers to C₂-C₁₆-alkenyl groups having a heteroaryl substituent

"C₂-C₁₂-alkynyl" refers to alkynyl groups preferably having from 2 to 12 carbon atoms and having at least 1-2 sites of alkynyl unsaturation. By analogy, "C₂-C₁₆-alkynyl" refers to alkynyl groups preferably having from 2 to 16 carbon atoms and having at least 1-2 sites of alkynyl unsaturation. Preferred alkynyl groups include ethynyl (-C≡CH), propargyl (-CH₂C≡CH), and the like.

"C₂-C₁₂-alkynyl aryl" refers to C₂-C₁₂-alkynyl groups having an aryl substituent, including phenylethynyl and the like. By analogy, "C₂-C₁₆-alkynyl aryl" refers to C₂-C₁₆-alkynyl groups having an aryl substituent.

"C₂-C₁₂-alkynyl heteroaryl" refers to C₂-C₁₂-alkynyl groups having a heteroaryl substituent, including 2-thienylethynyl and the like. By analogy, "C₂-C₁₆-alkynyl heteroaryl" refers to C₂-C₁₆-alkynyl groups having a heteroaryl substituent

"C₃-C₈-cycloalkyl" refers to a saturated carbocyclic group of from 3 to 8 carbon atoms having a single ring (*e.g.*, cyclohexyl) or multiple condensed rings (*e.g.*, norbornyl). Preferred cycloalkyl include cyclopentyl, cyclohexyl, norbornyl and the like.

"Heterocycloalkyl" refers to a C₃-C₈-cycloalkyl group according to the definition above, in which up to 3 carbon atoms are replaced by heteroatoms chosen from the group consisting of O, S, NR, R being defined as hydrogen, C₁₋₆ alkyl, alkoxy and the like. Preferred heterocycloalkyl include pyrrolidine, piperidine, piperazine, 1-methylpiperazine, morpholine, and the like.

"C₁-C₁₂-alkyl cycloalkyl" refers to C₁-C₁₂-alkyl groups having a cycloalkyl substituent, including cyclohexylmethyl, cyclopentylpropyl, and the like. By analogy, "C₁-C₁₆-alkyl cycloalkyl" refers to C₁-C₁₆-alkyl groups having a cycloalkyl substituent

"C₁-C₁₂-alkyl heterocycloalkyl" refers to C₁-C₁₂-alkyl groups having a heterocycloalkyl substituent, including 2-(1-pyrrolidinyl)ethyl, 4-morpholinylmethyl, (1-methyl-4-piperidinyl)methyl and the like. By analogy, "C₁-C₁₆-alkyl heterocycloalkyl" refers to C₁-C₁₆-alkyl groups having a heterocycloalkyl substituent

"Carboxy" refers to the group -C(O)OH.

"C₁-C₁₂-alkyl carboxy" refers to C₁-C₁₂-alkyl groups having a carboxy substituent, including 2-carboxyethyl and the like. By analogy, "C₁-C₁₆-alkyl carboxy" refers to C₁-C₁₆-alkyl groups having a carboxy substituent

"Acyl" refers to the group -C(O)R where R includes "C₁-C₁₂-alkyl", "C₁-C₈-alkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl".

"C₁-C₁₂-alkyl acyl" refers to C₁-C₁₂-alkyl groups having an acyl substituent, including 2-acetyethyl and the like. By analogy, "C₁-C₁₆-alkyl acyl" refers to C₁-C₁₆-alkyl groups having an acyl substituent

"Aryl acyl" refers to aryl groups having an acyl substituent, including 2-acetylphenyl and the like.

"Heteroaryl acyl" refers to heteroaryl groups having an acyl substituent, including 2-acetylpyridyl and the like.

"C₃-C₈-(hetero)cycloalkyl acyl" refers to 3 to 8 membered cycloalkyl or heterocycloalkyl groups having an acyl substituent.

"Acyloxy" refers to the group -OC(O)R where R includes H, "C₁-C₁₂-alkyl", "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl".

"C₁-C₁₂-alkyl acyloxy" refers to C₁-C₁₂-alkyl groups having an acyloxy substituent, including 2-(acetyloxy)ethyl and the like. By analogy, "C₁-C₁₆-alkyl acyloxy" refers to C₁-C₁₆-alkyl groups having an acyloxy substituent

"Alkoxy" refers to the group -O-R where R includes "C₁-C₁₂-alkyl", "C₁-C₈-alkyl", "C₁-C₆-alkyl" or "aryl" or "hetero-aryl" or "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl". Preferred alkoxy groups include by way of example, methoxy, ethoxy, phenoxy and the like.

"C₁-C₁₂-alkyl alkoxy" refers to C₁-C₁₂-alkyl groups having an alkoxy substituent, including 2-ethoxyethyl and the like. By analogy, "C₁-C₁₆-alkyl alkoxy" refers to C₁-C₁₆-

alkyl groups having an alkoxy substituent and "C₁-C₆-alkyl" refers to C₁-C₆-alkyl groups having an alkoxy substituent.

"Alkoxycarbonyl" refers to the group -C(O)OR where R includes H, "C₁-C₁₂-alkyl" or "aryl" or "heteroaryl" or "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl".

"C₁-C₁₂-alkyl alkoxycarbonyl" refers to C₁-C₁₂-alkyl groups having an alkoxycarbonyl substituent, including 2-(benzyloxycarbonyl)ethyl and the like. By analogy, "C₁-C₁₆-alkyl alkoxycarbonyl" refers to C₁-C₁₆-alkyl groups having an alkoxycarbonyl substituent

"Aminocarbonyl" refers to the group -C(O)NRR' where each R, R' includes independently hydrogen or C₁-C₁₂-alkyl or C₁-C₆-alkyl or aryl or heteroaryl or "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl hetero-aryl".

"C₁-C₁₂-alkyl aminocarbonyl" refers to C₁-C₁₂-alkyl groups having an aminocarbonyl substituent, including 2-(dimethylaminocarbonyl)ethyl and the like. By analogy, "C₁-C₁₆-alkyl aminocarbonyl" refers to C₁-C₁₆-alkyl groups having an aminocarbonyl substituent and "C₁-C₈-alkyl aminocarbonyl" refers to C₁-C₈-alkyl groups having an aminocarbonyl substituent, including 2-(dimethylaminocarbonyl)ethyl and the like.

"Acylamino" refers to the group -NRC(O)R' where each R, R' is independently hydrogen, "C₁-C₁₂-alkyl", "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl".

"C₁-C₁₂-alkyl acylamino" refers to C₁-C₁₂-alkyl groups having an acylamino substituent, including 2-(propionylamino)ethyl and the like. By analogy, "C₁-C₁₆-alkyl acylamino" refers to C₁-C₁₆-alkyl groups having an acylamino substituent

"Ureido" refers to the group -NRC(O)NR'R'' where each R, R', R'' is independently hydrogen, "C₁-C₁₂-alkyl", "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl", and where R' and R'', together with the nitrogen atom to which they are attached, can optionally form a 3-8-membered heterocycloalkyl ring.

"C₁-C₁₂-alkyl ureido" refers to C₁-C₁₂-alkyl groups having an ureido substituent, including 2-(N'-methylureido)ethyl and the like. By analogy, "C₁-C₁₆-alkyl ureido" refers to C₁-C₁₆-alkyl groups having an ureido substituent

"Carbamate" refers to the group -NRC(O)OR' where each R, R' is independently hydrogen, "C₁-C₁₂-alkyl", "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl".

"Amino" refers to the group -NRR' where each R, R' is independently hydrogen or "C₁-C₁₂-alkyl" or "aryl" or "heteroaryl" or "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", or "cycloalkyl", or "heterocycloalkyl", and where R and R', together with the nitrogen atom to which they are attached, can optionally form a 3-8-membered heterocycloalkyl ring.

"C₁-C₁₂-alkyl amino" refers to C₁-C₁₂-alkyl groups having an amino substituent, including 2-(1-pyrrolidinyl)ethyl and the like. By analogy, "C₁-C₁₆-alkyl amino" refers to C₁-C₁₆-alkyl groups having an amino substituent and "C₁-C₈-alkyl amino" refers to C₁-C₈-alkyl groups having an amino substituent

"Ammonium" refers to a positively charged group $\text{-N}^+\text{RR'R''}$, where each R, R', R'' is independently "C₁-C₁₂-alkyl" or "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", or "cycloalkyl", or "heterocycloalkyl", and where R and R', together with

the nitrogen atom to which they are attached, can optionally form a 3-8-membered heterocycloalkyl ring.

"C₁-C₁₂-alkyl ammonium" refers to C₁-C₁₂-alkyl groups having an ammonium substituent, including 2-(1-pyrrolidiny)ethyl and the like. By analogy, "C₁-C₁₆-alkyl ammonium" refers to C₁-C₁₆-alkyl groups having an ammonium substituent

"Halogen" refers to fluoro, chloro, bromo and iodo atoms.

"Sulfonyloxy" refers to a group -OSO₂-R wherein R is selected from H, "C₁-C₁₂-alkyl", "C₁-C₁₂-alkyl" substituted with halogens, *e.g.*, an -OSO₂-CF₃ group, "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl".

"C₁-C₁₂-alkyl sulfonyloxy" refers to C₁-C₁₂-alkyl groups having a sulfonyloxy substituent, including 2-(methylsulfonyloxy)ethyl and the like. By analogy, "C₁-C₁₆-alkyl sulfonyloxy" refers to C₁-C₁₆-alkyl groups having a sulfonyloxy substituent

"Sulfonyl" refers to group "-SO₂-R" wherein R is selected from H, "aryl", "heteroaryl", "C₁-C₁₂-alkyl", "C₁-C₁₂-alkyl" substituted with halogens, *e.g.*, an -SO₂-CF₃ group, "C₂-C₆-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl".

"C₁-C₁₂-alkyl sulfonyl" refers to C₁-C₁₂-alkyl groups having a sulfonyl substituent, including 2-(methylsulfonyl)ethyl and the like. By analogy, "C₁-C₁₆-alkyl sulfonyl" refers to C₁-C₁₆-alkyl groups having a sulfonyl substituent.

"Sulfinyl" refers to a group "-S(O)-R" wherein R is selected from H, "C₁-C₁₂-

alkyl", "C₁-C₆-alkyl" substituted with halogens, *e.g.*, an -SO-CF₃ group, "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl".

"C₁-C₁₂-alkyl sulfinyl" refers to C₁-C₁₂-alkyl groups having a sulfinyl substituent, including 2-(methylsulfinyl)ethyl and the like. By analogy, "C₁-C₁₆-alkyl sulfinyl" refers to C₁-C₁₆-alkyl groups having a sulfinyl substituent

"Sulfanyl" refers to groups -S-R where R includes H, "C₁-C₁₂-alkyl", "C₁-C₁₂-alkyl" substituted with halogens, *e.g.*, a -SO-CF₃ group, "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₆-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl". Preferred sulfanyl groups include methylsulfanyl, ethylsulfanyl, and the like.

"C₁-C₁₂-alkyl sulfanyl" refers to C₁-C₁₂-alkyl groups having a sulfanyl substituent, including 2-(ethylsulfanyl)ethyl and the like. By analogy, "C₁-C₁₆-alkyl sulfanyl" refers to C₁-C₁₆-alkyl groups having a sulfanyl substituent.

"Sulfonylamino" refers to a group -NRSO₂-R' where each R, R' includes independently hydrogen, "C₁-C₁₂-alkyl", "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl".

"C₁-C₁₂-alkyl sulfonylamino" refers to C₁-C₁₂-alkyl groups having a sulfonylamino substituent, including 2-(ethylsulfonylamino)ethyl and the like. By analogy, "C₁-C₁₆-alkyl sulfonylamino" refers to C₁-C₁₆-alkyl groups having a sulfonylamino substituent


"Aminosulfonyl" refers to a group $-\text{SO}_2\text{-NRR}'$ where each R, R' includes independently hydrogen, "C₁-C₁₂-alkyl", "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₁₂-alkyl aryl" or "C₁-C₁₂-alkyl heteroaryl", "C₂-C₁₂-alkenyl aryl", "C₂-C₁₂-alkenyl heteroaryl", "C₂-C₁₂-alkynyl aryl", "C₂-C₁₂-alkynylheteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl".

"C₁-C₁₂-alkyl aminosulfonyl" refers to C₁-C₁₂-alkyl groups having an aminosulfonyl substituent, including 2-(cyclohexylaminosulfonyl)ethyl and the like. By analogy, "C₁-C₁₆-alkyl aminosulfonyl" refers to C₁-C₁₆-alkyl groups having an aminosulfonyl substituent.


"Substituted or unsubstituted" or "optionally substituted: Unless otherwise constrained by the definition of the individual substituent, the above set out groups, like "alkyl", "alkenyl", "alkynyl", "aryl" and "heteroaryl" etc. groups can optionally be substituted with from 1 to 5 substituents selected from the group consisting of "C₁-C₁₂-alkyl", "C₂-C₁₂-alkenyl", "C₂-C₁₂-alkynyl", "cycloalkyl", "heterocycloalkyl", "C₁-C₁₂-alkyl aryl", "C₁-C₁₂-alkyl heteroaryl", "C₁-C₁₂-alkyl cycloalkyl", "C₁-C₁₂-alkyl heterocycloalkyl", "amino", "ammonium", "acyl", "acyloxy", "acylamino", "aminocarbonyl", "alkoxycarbonyl", "ureido", "aryl", "carbamate", "heteroaryl", "sulfinyl", "sulfonyl", "alkoxy", "sulfanyl", "halogen", "carboxy", trihalomethyl, cyano, hydroxy, mercapto, nitro, and the like. Alternatively said substitution could also comprise situations where neighbouring substituents have undergone ring closure, notably when vicinal functional substituents are involved, thus forming, e.g., lactams, lactone, cyclic anhydrides, but also acetals, thioacetals, amins formed by ring closure for instance in an effort to obtain a protective group.

Specifically, preferred compounds of the invention include the following compounds, and pharmaceutically acceptable salts of these compounds, with the compound name set forth above the depicted compound structure. The structurally depicted compounds generally show preferred stereoisomers.


4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridinyl-3-yl-1H-benzoimidazol-5-yl)-amide




4-pentanoyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl) amide




{[3-(9-ethyl-9H-carbazol-3-yl)carbamoyl]-4-(thiophene-2-sulfonyl)-piperazine-1-carbonyl]-amino}acetic acid ethyl ester




4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 1-ethylamide 3-[(9-ethyl-9H-carbazol-3-yl)amide]




4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)amide]-1-pentylamide




4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl) amide




4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(2-methoxy-ethyl)-amide]




4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 1-pentylamide 3-[(3-pyridin-4-yl-phenyl)-amide]



4-(1-methyl-1H-imidazole-4-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-pentylamide




4-(dimethylsulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-pentylamide




4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)-amide


24




4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl)-amide




4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(3-imidazol-1-yl-propyl)-amide]




4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(3-imidazol-1-yl)-amide]




4-pentyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl) amide




4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl) amide



4-(3-methylsulfonyl-propyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl)-amide




4-(4-ethyl-furan-3-ylmethyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl)-amide




3-(9-ethyl-9H-carbazol-3-yl)-4-(thiophene-2-sulfonyl)-piperazin-1-yl] acetic acid ethyl ester


27




1-benzenesulfonyl-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide




4-pentyl-1-phenylsulfonyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide




4-hexyl-1-thiophene-2-sulfonyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide



1-(4-fluoro-benzenesulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide




1-(2-fluoro-benzenesulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide




4-octyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide

29




4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzoimidazol-5-yl) amide




/


1-(butane-1-sulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide




4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide



4-(3-methylsulfonyl-propyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide




4-octyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide



Compounds of the invention can be readily prepared. For instance, a suitable reaction sequence to provide compounds of the invention is set forth in the following Schemes 1A and 1B:

32

Scheme 1A




As exemplified in the above Scheme 1A of the ring nitrogens of a 2-carboxylic acid piperazine 1a (1) can be protected e.g. with FMOC chloride (9-fluorenylmethoxycarbonyl chloride) to lead a compound 2a (2). The unprotected ring nitrogen of compound 2a can be reacted with a sulfonyl reagent to introduce the -SO₂R⁴ substituent of Formulae I to III, leading to sulfonyl derivatives of formula 3a such as for example an optionally substituted thiophene sulfonyl compound 3 as shown in Scheme 1B below.

The 2-carboxy group then can be functionalized to provide the carboxamide moiety, suitably by reaction with a primary or secondary amine of formula HNR¹R², where the amine reagent substituents provide R¹ and R² groups as defined in Formulae I to III above, such as to provide amide derivative of formula 4a, such as for example compound 4 as shown in Scheme 1B below.

The protected ring nitrogen then can be deprotected e.g. in the presence of base such as diethyl amine or other organic base, to provide a compound of formula 5a, for example a compound 5 of Scheme 1B below. The deprotected ring nitrogen then may be

substituted as desired to provide an R^3 substituent, as defined in Formulae I to III above to lead to a compound of formula 6a, e.g. reacted with an optionally substituted alkyl chloride, optionally substituted benzyl chloride and the like to provide the corresponding R^3 moiety, such as compound 6 in the Scheme 1B below.

Scheme 1B



If the above set out general synthetic methods are not applicable for obtaining certain compounds of formula I, suitable methods of preparation known by a person skilled in the art should be used.

A particularly preferred reaction route to provide compounds of the invention is set forth in Example 1, which follows.

As discussed above, the present invention includes methods for treating infertility in male and female mammals, such as primates, particularly humans. Compounds of the invention will be useful for treatment of infertility conditions currently treated with FSH and/or LH, including female infertility and male spermatogenesis disorders.

The therapeutic methods of the invention generally comprise administration of an effective amount of one or more compounds of the invention to a subject including a mammal, such as a primate, especially a human, in need of such treatment.

Typical candidates for treatment in accordance with the methods of the invention persons suffering from or suspected of suffering from infertility. See the *Merck Manual*, vol. 2, pages 12-17 (16th ed.) for identification of patients suffering from or suspected of infertility, which in the case of humans, can include failure to conceive within one year of unprotected intercourse.

The treatment methods of the invention may be particularly beneficial for female mammals suffering from an ovulatory disorder. Additionally, compounds of the invention can be administered to females undergoing assisted reproductive treatments such as in-vitro fertilization, e.g. to simulate follicular development and maturation. Compounds of the invention also can be administered to males to facilitate adequate spermatogenesis.

The treatment methods of the invention also will be useful for treatment of infertility in mammals other than humans, such as horses and livestock e.g. cattle, sheep, cows and the like.

Compounds of the invention may be administered as a "cocktail" formulation, i.e. coordinated administration of one or more compounds of the invention together with one or more other active therapeutics, particularly one or more other known fertility agents. For instance, one or more compounds of the invention may be administered in coordination with a regime of Follicle Stimulating Hormone and/or Leutinizing Hormone such as Gonal-F, Metrodin HP or Pergonal, or other agent such as clomephine citrate.

The compounds of this invention can be administered by a variety of routes, such as orally or by injection, e.g., intramuscular, intraperitoneal, subcutaneous or intravenous injection, or topically e.g. transdermally, vaginally and the like. Compounds of the invention may be suitably administered to a subject in the protonated and water-soluble

form, e.g., as a pharmaceutically acceptable salt of an organic or inorganic acid, e.g., hydrochloride, sulfate, hemi-sulfate, phosphate, nitrate, acetate, oxalate, citrate, maleate, mesylate, etc.

Compounds of the invention can be employed, either alone or in combination with one or more other therapeutic agents as discussed above, as a pharmaceutical composition in mixture with conventional excipient, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for oral, parenteral, enteral or topical application which do not deleteriously react with the active compounds and are not deleterious to the recipient thereof. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds.

For oral administration, pharmaceutical compositions containing one or more substituted piperazine compounds of the invention may be formulated as e.g. tablets, troches, lozenges, aqueous or oil suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups, elixirs and the like. Typically, suitable are tablets, dragees or capsules having talc and/or carbohydrate carrier binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein the active component is protected with differentially degradable coatings, e.g., by microencapsulation, multiple coatings, etc.

For parenteral application, e.g., sub-cutaneous, intraperitoneal or intramuscular, particularly suitable are solutions, preferably oil or aqueous solutions, as well as

suspensions, emulsions, or implants, including suppositories. Ampules are convenient unit dosages.

For topical applications, formulations may be prepared in a topical ointment or cream containing one or more compounds of the invention. When formulated as an ointment, one or more compounds of the invention suitably may be employed with either a paraffinic or a water-miscible base. The one or more compounds also may be formulated with an oil-in-water cream base. Other suitable topical formulations include e.g. lozenges and dermal patches.

It will be appreciated that the actual preferred amounts of active compounds used in a given therapy will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, the particular site of administration, etc. Optimal administration rates for a given protocol of administration can be readily ascertained by those skilled in the art using conventional dosage determination tests conducted with regard to the foregoing guidelines. See also Remington's Pharmaceutical Sciences. In general, a suitable effective dose of one or more compounds of the invention, particularly when using the more potent compound(s) of the invention, will be in the range of from 0.01 to 100 milligrams per kilogram of bodyweight of recipient per day, preferably in the range of from 0.01 to 20 milligrams per kilogram bodyweight of recipient per day, more preferably in the range of 0.05 to 4 milligrams per kilogram bodyweight of recipient per day. The desired dose is suitably administered once daily, or several sub-doses, e.g. 2 to 4 sub-doses, are administered at appropriate intervals throughout the day, or other appropriate schedule. Such sub-doses may be administered as unit dosage forms, e.g., containing from 0.05 to 10 milligrams of compound(s) of the invention, per unit dosage.

The entire text of all documents cited herein are incorporated by reference herein. The following non-limiting examples are illustrative of the invention.


The following abbreviations refer respectively to the definitions below:

h (hour), g (gram), mg (milligram), min (minute), mM (millimolar), mL (milliliter). cAmP (Cyclic Adenosine Monophosphate); CHO (Chinese Hamster Ovary); DMF (dimethylformamide); DMSO (Dimethyl Sulfoxide); FSH (Follicle Stimulating Hormone); Fmoc (9-fluorenylmethoxycarbonyl); Gs (G-proteins); HATU (0-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate); HTS (High-Troughput Screening); LDR (Log-Dose Response); PDE (Phosphodiesterase); RT (Room Temperature); TSH (Tyroid Stimulating Hormone).

Example 1: Synthesis of 4-Hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl)-amide

The title compound was prepared as outlined in the following Scheme 2.

Scheme 2



Part 1: Synthesis of Piperazine-1,3(R)-dicarboxylic acid-1-(9H-fluoren-9-ylmethyl) ester (Compound 2 in above Scheme 2).

Piperazine-2(R)-carboxylic acid dihydrochloride (5.95g) was added to 50 mL dioxane/water (1:1). The solution was cooled in an ice bath and aqueous sodium hydroxide (10% w/v) solution was added drop-wise to adjust the pH to 9~10. At this pH, piperazine-2-carboxylic acid dissolved completely. Then, Fmoc Chloride (2.46 g) in dioxane was added drop-wise to the ice-cooled solution of piperazine and the reaction mixture was stirred from 0°C to RT over 5-6 hr. The mixture was poured into 800 ml of ice water and washed once with ether. Then the aqueous phase was acidified to pH 2-3 with 2N of HCl which precipitated a white solid. This solid was filtered and dried in

vacuo to obtain 2.46g of a white powder. MS (ESI pos.): 353 (M+H).

Part 2: Synthesis of 4-(Thiophene-2-sulfonyl)-piperazine-1,3(R)-dicarboxylic acid 1-(9H-fluoren-9-ylmethyl) ester (Compound 3 in above Scheme 2).

4-Fmoc protected piperazine-2(R)-carboxylic acid (352 mg) was dissolved in 10 mL anhydrous methylene chloride. The solution was cooled in an ice bath and 0.35 mL di-isopropyl ethyl amine was added. Then, 183 mg of 2-thiophene sulfonyl chloride was added and the reaction mixture was stirred at 0°C for 5hr. The mixture was diluted with dichloromethane, washed with 0.5N aqueous hydrochloric acid and brine. The solvent was evaporated from the organic layer to leave a yellowish solid which was dried in vacuo. This crude material was used as such for the next step without purification.

Part 3: Synthesis of 3-(1-Ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-ylcarbamoyl)-4-(thiophene-2-sulfonyl)-piperazine-1-carboxylic acid 9H-fluoren-9-ylmethyl ester (Compound 4 in above Scheme 2).

The crude material from the previous step was dissolved in anhydrous DMF and to this were added 238 mg of 1-Ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-ylamine followed by 258 mg of di-isopropyl ethyl amine and 380 mg of HATU. The mixture was stirred for 5hr upon which the crude mixture was diluted with ethyl acetate and the organic layer was successively washed with saturated sodium bicarbonate, 0.5N HCl and brine. The organic layer was evaporated to dryness and the product was isolated by chromatography over silica gel (0 to 5% methanol in ethyl acetate). MS (ESI pos.): 719 (M+H).

Part 4: Deprotection of Fmoc (9H-fluoren-9-ylmethyl ester) group:

The Fmoc-protected piperazine derivative from step 3 was dissolved in 5 mL dichloromethane. To this were slowly added 5 mL of diethyl amine and the resulting mixture was stirred for 1.5 hr at ambient temperature. The excess reagent was evaporated in vacuo and the crude product was purified by chromatography over silica gel (0-25% methanol in ethyl acetate) to afford the piperazine 5. MS (ESI pos.): 497 (M+H)

Part 5: Synthesis of 4-Hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl)-amide (Compound **6** in above Scheme 2).

The piperazine **5** (200 mg) as prepared in Part 4 above was dissolved in dichloromethane containing 5% acetic acid and 144 microliters of n-hexanal were added. After stirring for 16 hours, 500 mg of sodium cyano borohydride was added and the reaction was stirred for 5 hr at room temperature. The crude mixture was washed with saturated sodium bicarbonate, brine and dried. Purification over silica gel (0-5% methanol in ethyl acetate) afforded 126 mg of the desired product **6**. MS (ESI pos.): 581 (M+H)

Example 2: FSH Assay Method

Compounds of the invention were tested in the following assay.

Assay Overview.

All compounds of the invention were stored in 96-well deepwell plates in DMSO at a nominal concentration of 10 μ M. Compounds of the invention were screened for agonist activity at the FSH receptor using the recombinant FSH receptor stably transfected and expressed in Chinese Hamster Ovary Cells (CHO cells) essentially as described in the work by Kelton, et al. (Molecular and Cellular Endocrinology, 1992, 89, 141-151). Since the FSH receptor is known to act via a G-protein (Gs) to activate adenylyl cyclase and hence raise intracellular levels of cAMP the high throughput screening (HTS) assay used a gene reporter system consisting of the cAMP response element coupled upstream to the reporter gene which, in this case was the enzyme luciferase. Thus an agonist at the FSH receptor increases cAMP in the cell which results in activation of the CREB (cAMP) response element binding protein). The substrate for the enzyme (Packard Instrument Company 800 Research Parkway, Meriden, CT 06450, USA) was added to the cells after appropriate incubation with the compounds or FSH and the amount of luciferase expressed was measured by quantitating the luminescence produced by the enzyme using a TopCount scintillation/luminescence counter running in single photon counting mode. A compound which acts as an agonist at the receptor should produce light from the treated cells in proportion to its concentration within the incubation. Luminescence should be saturable at high concentrations of the compound.

Protocol.

The compounds of the invention, in deepwell plates (Master plates) as indicated above were loaded on the robotic deck along with the appropriate number of assay plates and daughter plates. A 10 μ l aliquot from each master plate was transferred to the corresponding daughter plate and 90 μ l of DME/F12 was added and mixed within each well. 20 μ l was then removed from the daughter plate and dispensed into the assay plate. After addition of an aliquot of FSH (equivalent to an EC100 response for this hormone to each of three wells on the plate, 80 μ l of media (DME/F 12 \pm 2% serum) and 100 μ l aliquot of cells (4×10^5 /ml in the same media) were added and the plate incubated at 37°C for 3 hrs 30 min. At this time the plate was removed from the incubator and media in each well was aspirated and the cells adhering to the bottom of the plate washed with 300 μ l PBS containing 1mM Ca²⁺ and 1mM Mg²⁺. The PBS was aspirated and 100 μ l PBS added to each well. 100 μ l of Lucite (prepared as described by the manufacturer) was added to each well and the plate was shaken gently for 40 seconds prior to placement in the TopCount plate reader. After allowing 3.5 minutes for the plate to dark-adapt within the machine, the amount of luminescence generated was quantitated using Single Photon Counting mode. The data was transmitted electronically from the TopCount to the robot processing computer terminal and was renamed with an ID corresponding to the original master plate ID. Data was evaluated using an Excel macro and compounds showing activity comparable to that produced by an ED100 of FSH itself were further analyzed in the same assay at differing concentrations. LDR (log-dose-response) curves were generated for these compounds in CHO cells containing the FSH receptor and these curves were also compared with those in either cells expressing a different Gs linked receptor or in cells lacking any transfected receptor. Results are set forth in the following Table 1.

Table 1

Tested Compound	ED ₅₀ (nM)
4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridinyl-3-yl-1H-benzimidazol-5-yl)-amide	40
4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)amide]-1-pentylamide	63
4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 1-ethylamide 3-[(9-ethyl-9H-carbazol-3-yl)amide]	877
{[3-(9-ethyl-9H-carbazol-3-yl)carbamoyl]-4-(thiophene-2-sulfonyl)-piperazine-1-carbonyl]-amino} acetic acid ethyl ester	408
4-pentanoyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl) amide	402
4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl) amide	380
4-dimethylsulfonyl-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)amide] 1-pentylamide	403
4-(1-methyl-1H-imidazole-4-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-pentylamide	3160
4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 1-pentylamide 3-[(3-pyridin-4-yl-phenyl)-amide]	5000
4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[[2-(1H-imidazol-4-yl)-ethyl]-amide]	6250

Table 1 (continued)

Tested Compound	ED ₅₀ (nM)
4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)-amide	978
4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl)-amide	94
4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(3-imidazol-1-yl-propyl)-amide]	6470
4-pentyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)amide	220
4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)amide	1200
4-(3-methylsulfonyl-propyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl)-amide	1700
4-(4-ethyl-furan-3-ylmethyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl)-amide	1600
3-(9-ethyl-9H-carbazol-3-ylcarbamoyl)-4-(thiophene-2-sulfonyl)-piperazin-1-yl] acetic acid ethyl ester	610
1-benzenesulfonyl-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide	150
4-pentyl-1-thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide	150
4-hexyl-1-thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide	180
1-(4-fluoro-benzenesulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide	167


Table 1 (continued)

Tested Compound	ED ₅₀ (nM)
1-(2-fluoro-benzenesulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide	74
4-octyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzoimidazol-5-yl) amide	54
4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzoimidazol-5-yl) amide	53
1-dimethylsulfamoyl-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide	242
1-(butane-1-sulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide	170
4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzoimidazol-5-yl) amide	111
4-(3-methylsulfanyl-propyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzoimidazol-5-yl) amide	14000
4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(2-methoxy-ethyl)-amide]	1610
4-octyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide	13

The invention has been described in detail with reference to preferred embodiments thereof. However, it will be appreciated that those skilled in the art, upon consideration of this disclosure, may make modifications and improvements within the spirit and scope of the invention.

What is claimed is:

1. A method for treating infertility in a mammal, comprising administering to a mammal suspected of infertility a therapeutically effective amount of a compound of Formula I:



I

wherein R¹ and R² are independently selected from the group comprising or consisting of hydrogen, C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₂-C₁₂-alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group, an acyl moiety, C₁-C₁₂-alkyl aryl, C₁-C₁₂-alkyl heteroaryl, C₂-C₁₂-alkenyl aryl, C₂-C₁₂-alkenyl heteroaryl, C₂-C₁₂-alkynyl aryl, C₂-C₁₂-alkynyl heteroaryl, C₁-C₁₂-alkyl cycloalkyl, C₁-C₁₂-alkyl heterocycloalkyl, C₂-C₁₂-alkenyl cycloalkyl, C₂-C₁₂-alkenyl heterocycloalkyl, C₂-C₁₂-alkynyl cycloalkyl, C₂-C₁₂-alkynyl heterocycloalkyl, alkoxycarbonyl, aminocarbonyl, C₁-C₁₂-alkyl carboxy, C₁-C₁₂-alkyl acyl, aryl acyl, heteroaryl acyl, C₃-C₈-(hetero)cycloalkyl acyl, C₁-C₁₂-alkyl acyloxy, C₁-C₁₂-alkyl alkoxy, C₁-C₁₂-alkyl alkoxycarbonyl, C₁-C₁₂-alkyl aminocarbonyl, C₁-C₁₂-alkyl acylamino, acylamino, C₁-C₁₂-alkyl ureido, C₁-C₁₂-alkyl carbamate, C₁-C₁₂-alkyl amino, C₁-C₁₂-alkyl ammonium, C₁-C₁₂-alkyl sulfonyloxy, C₁-C₁₂-alkyl sulfonyl, C₁-C₁₂-alkyl sulfinyl, C₁-C₁₂-alkyl sulfanyl, C₁-C₁₂-alkyl sulfonylamino, or C₁-C₁₂-alkyl aminosulfonyl;


R³ is C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₂-C₁₆-alkynyl, wherein said alkyl, alkenyl, alkynyl

chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group, an acyl moiety, C₁-C₁₆-alkyl aryl, C₁-C₁₆-alkyl heteroaryl, C₂-C₁₆-alkenyl aryl, C₂-C₁₆-alkenyl heteroaryl, C₂-C₁₆-alkynyl aryl, C₂-C₁₆-alkynyl heteroaryl, C₁-C₁₆-alkyl cycloalkyl, C₁-C₁₆-alkyl heterocycloalkyl, C₂-C₁₆-alkenyl cycloalkyl, C₂-C₁₆-alkenyl heterocycloalkyl, C₂-C₁₆-alkynyl cycloalkyl, C₂-C₁₆-alkynyl heterocycloalkyl, alkoxycarbonyl, aminocarbonyl, C₁-C₁₆-alkyl carboxy, C₁-C₁₆-alkyl acyl, aryl acyl, heteroaryl acyl, C₃-C₈-(hetero)cycloalkyl acyl, C₁-C₁₆-alkyl acyloxy, C₁-C₁₆-alkyl alkoxy, C₁-C₁₆-alkyl alkoxycarbonyl, C₁-C₁₆-alkyl aminocarbonyl, C₁-C₁₆-alkyl acylamino, acylamino, C₁-C₁₆-alkyl ureido, C₁-C₁₆-alkyl carbamate, C₁-C₁₆-alkyl amino, C₁-C₁₆-alkyl ammonium, C₁-C₁₆-alkyl sulfonyloxy, C₁-C₁₆-alkyl sulfonyl, C₁-C₁₆-alkyl sulfinyl, C₁-C₁₆-alkyl sulfanyl, C₁-C₁₆-alkyl sulfonylamino, or C₁-C₁₆-alkyl aminosulfonyl;

R⁴ is C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₂-C₁₂-alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group, or amino; and pharmaceutically acceptable salts thereof.


2. A method according to claim 1, wherein the compound of Formula I is such as R¹ is H.
3. A method according to claims 1 or 2, wherein the compound of Formula I is such as R² is selected from aryl, heteroaryl, 3-8 membered cycloalkyl and heterocycloalkyl.
4. A method according to any of the preceding claims, wherein the compound of Formula I is such as R⁴ is selected from C₁-C₆-alkyl, amino, aryl, heteroaryl, 3-8-membered cycloalkyl and heterocycloalkyl.

5. A method of treatment according to any of the preceding claims, wherein the compound of Formula I is such as R^1 is H; R^2 is aryl; R^3 is selected from C_1 - C_8 -alkyl, C_1 - C_8 -acyl amino and C_1 - C_8 -alkyl acyl and R^4 is selected from C_1 - C_6 -alkyl, amino, aryl and heteroaryl.
6. A method according to any of the preceding claims wherein the compound has the following Formula II:

**II**

wherein R^1 , R^2 and R^3 are the same as defined above in Formula I; each R^5 is independently halogen, hydroxy or the same as defined for R^1 ; m is an integer of from 0 to 4; and pharmaceutically acceptable salts thereof.

7. A method of claim 1 wherein the compound has the following Formula III:

**III**

wherein R^1 , R^2 and R^3 are each the same as defined above in Formula I; each R^5 is independently halogen, hydroxy or the same as defined for R^1 ; m is an integer of from 0 to 4; and pharmaceutically acceptable salts thereof.

8. A method according to any of the preceding claims wherein R^1 is hydrogen and R^2 is other than hydrogen.
9. A method according to any of the preceding claims wherein R^2 is aryl or heteroaryl.
10. A method according to any of the preceding claims wherein R^3 is alkyl having five or more carbon atoms.
11. A method of claim 10 wherein R^3 is an n-alkyl group.
12. A method of claim 1 wherein R^4 is optionally substituted alkyl, aryl heteroaryl.
13. A method of any one of claims 1 through 12 wherein R^2 comprises a carbazolyl, tetrahydro-beta-carbolinyl or benzimidazolyl moiety.
14. A method according to any of the preceding claims wherein the compound of formula I is selected from the following group:
 - 4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridinyl-3-yl-1H-benzimidazol-5-yl)-amide;
 - 4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)amide]-1-pentylamide;
 - 4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 1-ethylamide 3-[(9-ethyl-9H-carbazol-3-yl)amide];
 - {[3-(9-ethyl-9H-carbazol-3-yl)carbonyl]-4-(thiophene-2-sulfonyl)-piperazine-1-carbonyl]-amino}acetic acid ethyl ester;
 - 4-pentanoyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl) amide;

4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl) amide;

4-dimethylsulfamoyl-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)amide] 1-pentylamide;

4-(1-methyl-1H-imidazole-4-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-pentylamide;

4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 1-pentylamide 3-[(3-pyridin-4-yl-phenyl)-amide];

4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[[2-(1H-imidazol-4-yl)-ethyl]-amide];

4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)-amide;

4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl)-amide;

4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(3-imidazol-1-yl-propyl)-amide];

4-pentyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)amide;

4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)amide;

4-(3-methylsulfanyl-propyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl)-amide;

4-(4-ethyl-furan-3-ylmethyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl)-amide;

3-(9-ethyl-9H-carbazol-3-ylcarbamoyl)-4-(thiophene-2-sulfonyl)-piperazin-1-yl] acetic acid ethyl ester;

1-benzenesulfonyl-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;

4-pentyl-1-thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;

4-hexyl-1-thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;

1-(4-fluoro-benzenesulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 1-(2-fluoro-benzenesulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 4-octyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide;
 4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide;
 1-dimethylsulfamoyl-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 1-(butane-1-sulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide;
 4-(3-methylsulfanyl-propyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide;
 4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(2-methoxy-ethyl)-amide];
 4-octyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide; and pharmaceutically acceptable salts thereof.

15. A method for treatment of a subject suffering from or susceptible to a disease or disorder associated with phosphodiesterase PDE4, adenosine transporters, or prostanoid receptors, comprising administering to the mammal a therapeutically effective amount of a compound of any one of claims 1 to 14.
16. A method of any one of claims 1 through 15 wherein the mammal is a human.
17. A method of any one of claims 1 through 16 wherein the mammal is a female.
18. A method of claim 17 wherein the mammal is suffering from an ovulatory

disorder.


19. A method of claim 17 wherein the mammal is being treated with an assisted reproduction procedure.

20. A method of claim 17 wherein the mammal is undergoing in-vitro fertilization.

21. A method of any one of claims 1 through 16 wherein the mammal is a male.

22. A method of any one of claims 21 wherein the mammal is a male suffering from a spermatogenesis disorder.

23. A compound according to Formula I :



I

wherein R¹ is H;


R² is selected from aryl, heteroaryl, 3-8-membered cycloalkyl and heterocycloalkyl;

R³ is selected from C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₂-C₁₆-alkynyl, monocyclic aryl, monocyclic heteroaryl, 3-8-membered monocyclic cycloalkyl, monocyclic heterocycloalkyl, acyl, C₁-C₁₆-alkyl aryl, C₁-C₁₆-alkyl heteroaryl, C₂-C₁₆-alkenyl aryl, C₂-C₁₆-alkenyl heteroaryl, C₂-C₁₆-alkynyl aryl, C₂-C₁₆-alkynyl heteroaryl, C₁-C₁₆-alkyl cycloalkyl, C₁-C₁₆-alkyl heterocycloalkyl, C₂-C₁₆-alkenyl cycloalkyl, C₂-C₁₆-alkenyl heterocycloalkyl, C₂-C₁₆-alkynyl cycloalkyl, C₂-C₁₆-alkynyl heterocycloalkyl, alkoxycarbonyl, aminocarbonyl, C₁-C₁₆-alkyl carboxy, C₁-C₁₆-alkyl acyl, aryl acyl, heteroaryl acyl, C₃-C₈-(hetero)cycloalkyl acyl, C₁-C₁₆-alkyl acyloxy, C₁-C₁₆-alkyl alkoxy, C₁-C₁₆-alkyl alkoxycarbonyl, C₁-C₁₆-alkyl aminocarbonyl, C₁-C₁₆-alkyl

acylamino, acylamino, C₁-C₁₆-alkyl sulfinyl, C₁-C₁₆-alkyl sulfanyl, C₁-C₁₆-alkyl ureido, C₁-C₁₆-alkyl carbamate, C₁-C₁₆-alkyl amino and C₁-C₁₆-alkyl ammonium; R⁴ is selected from C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₂-C₁₂-alkynyl, aryl, heteroaryl, 3-8-membered cycloalkyl, heterocycloalkyl, and amino; and pharmaceutically acceptable salts thereof.

24. A compound according to claim 23 wherein R² is selected from aryl, heteroaryl, 3-8 membered cycloalkyl and heterocycloalkyl.
25. A compound according to claims 23 or 24, wherein R⁴ is selected from C₁-C₆-alkyl, C₁-C₆-alkyl amino, aryl, heteroaryl, 3-8-membered cycloalkyl and heterocycloalkyl.
26. A compound according to any one of claims 23 through 25, wherein R² is aryl; R³ is selected from C₁-C₈-alkyl, C₁-C₈-acyl amino and C₁-C₈-alkyl acyl and R⁴ is selected from C₁-C₆-alkyl, amino, aryl and heteroaryl.
27. A compound according to any one of claims 23 through 26, wherein R² is fused phenyl
28. A compound according to any one of claims 23 through 27, wherein R⁴ is thienyl
29. A compound according to any one of claims 23 through 28 having the following Formula II:

53



II

wherein R^1 and R^2 are independently selected from the group comprising or consisting of hydrogen, C_1 - C_{12} -alkyl, C_2 - C_{12} -alkenyl, C_2 - C_{12} -alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group, an acyl moiety, C_1 - C_{12} -alkyl aryl, C_1 - C_{12} -alkyl heteroaryl, C_2 - C_{12} -alkenyl aryl, C_2 - C_{12} -alkenyl heteroaryl, C_2 - C_{12} -alkynyl aryl, C_2 - C_{12} -alkynyl heteroaryl, C_1 - C_{12} -alkyl cycloalkyl, C_1 - C_{12} -alkyl heterocycloalkyl, C_2 - C_{12} -alkenyl cycloalkyl, C_2 - C_{12} -alkenyl heterocycloalkyl, C_2 - C_{12} -alkynyl cycloalkyl, C_2 - C_{12} -alkynyl heterocycloalkyl, alkoxycarbonyl, aminocarbonyl, C_1 - C_{12} -alkyl carboxy, C_1 - C_{12} -alkyl acyl, aryl acyl, heteroaryl acyl, C_3 - C_8 -(hetero)cycloalkyl acyl, C_1 - C_{12} -alkyl acyloxy, C_1 - C_{12} -alkyl alkoxy, C_1 - C_{12} -alkyl alkoxycarbonyl, C_1 - C_{12} -alkyl aminocarbonyl, C_1 - C_{12} -alkyl acylamino, acylamino, C_1 - C_{12} -alkyl ureido, C_1 - C_{12} -alkyl carbamate, C_1 - C_{12} -alkyl amino, C_1 - C_{12} -alkyl ammonium, C_1 - C_{12} -alkyl sulfonyloxy, C_1 - C_{12} -alkyl sulfonyl, C_1 - C_{12} -alkyl sulfinyl, C_1 - C_{12} -alkyl sulfanyl, C_1 - C_{12} -alkyl sulfonylamino, or C_1 - C_{12} -alkyl aminosulfonyl;

R^3 is C_1 - C_{16} -alkyl, C_2 - C_{16} -alkenyl, C_2 - C_{16} -alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused

with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group, an acyl moiety, C₁-C₁₆-alkyl aryl, C₁-C₁₆-alkyl heteroaryl, C₂-C₁₆-alkenyl aryl, C₂-C₁₆-alkenyl heteroaryl, C₂-C₁₆-alkynyl aryl, C₂-C₁₆-alkynyl heteroaryl, C₁-C₁₆-alkyl cycloalkyl, C₁-C₁₆-alkyl heterocycloalkyl, C₂-C₁₆-alkenyl cycloalkyl, C₂-C₁₆-alkenyl heterocycloalkyl, C₂-C₁₆-alkynyl cycloalkyl, C₂-C₁₆-alkynyl heterocycloalkyl, alkoxycarbonyl, aminocarbonyl, C₁-C₁₆-alkyl carboxy, C₁-C₁₆-alkyl acyl, aryl acyl, heteroaryl acyl, C₃-C₈-(hetero)cycloalkyl acyl, C₁-C₁₆-alkyl acyloxy, C₁-C₁₆-alkyl alkoxy, C₁-C₁₆-alkyl alkoxycarbonyl, C₁-C₁₆-alkyl aminocarbonyl, C₁-C₁₆-alkyl acylamino, acylamino, C₁-C₁₆-alkyl ureido, C₁-C₁₆-alkyl carbamate, C₁-C₁₆-alkyl amino, C₁-C₁₆-alkyl ammonium, C₁-C₁₆-alkyl sulfonyloxy, C₁-C₁₆-alkyl sulfonyl, C₁-C₁₆-alkyl sulfinyl, C₁-C₁₆-alkyl sulfanyl, C₁-C₁₆-alkyl sulfonylamino, or C₁-C₁₆-alkyl aminosulfonyl;

R⁴ is C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₂-C₁₂-alkynyl, wherein said alkyl, alkenyl, alkynyl chains may be interrupted by a heteroatom selected from N, O or S, aryl, heteroaryl, saturated or unsaturated 3-8-membered cycloalkyl, heterocycloalkyl, wherein said cycloalkyl, heterocycloalkyl, aryl or heteroaryl groups may be fused with 1-2 further cycloalkyl, heterocycloalkyl, aryl or heteroaryl group;

R⁵ is independently halogen, hydroxy or the same as defined for R¹; m is an integer of from 0 to 4.

30. A compound according to any of the claims 23 to 29 wherein R² comprises a carbazolyl, tetrahydro-beta-carbolinyl or a benzimidazolyl moiety.

31. A compound according to any of the claims 23 to 30 that is selected from the following group:

4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridinyl-3-yl-1H-benzimidazol-5-yl)-amide);

4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)amide]-1-pentylamide;

4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 1-ethylamide 3-[(9-ethyl-9H-carbazol-3-yl)amide];

{[3-(9-ethyl-9H-carbazol-3-ylcarbamoyl)-4-(thiophene-2-sulfonyl)-piperazine-1-carbonyl]-amino} acetic acid ethyl ester;

4-pentanoyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl) amide;

4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl) amide;

4-dimethylsulfamoyl-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)amide] 1-pentylamide;

4-(1-methyl-1H-imidazole-4-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-pentylamide;

4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 1-pentylamide 3-[(3-pyridin-4-yl-phenyl)-amide];

4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[[2-(1H-imidazol-4-yl)-ethyl]-amide];

4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)-amide;

4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl)-amide;

4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(3-imidazol-1-yl-propyl)-amide];

4-pentyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)amide;

4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-oxo-2,3,4,9-tetrahydro-1H-beta-carbolin-6-yl)amide;

4-(3-methylsulfanyl-propyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl)-amide;

4-(4-ethyl-furan-3-ylmethyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (9-ethyl-9H-carbazol-3-yl)-amide;

3-(9-ethyl-9H-carbazol-3-ylcarbamoyl)-4-(thiophene-2-sulfonyl)-piperazine-1-yl] acetic acid ethyl ester;

1-benzenesulfonyl-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzoimidazol-5-yl) amide;

4-pentyl-1-thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 4-hexyl-1-thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 1-(4-fluoro-benzenesulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 1-(2-fluoro-benzenesulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 4-octyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide;
 4-heptyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide;
 1-dimethylsulfamoyl-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 1-(butane-1-sulfonyl)-4-hexyl-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide;
 4-hexyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide;
 4-(3-methylsulfanyl-propyl)-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid (1-ethyl-2-pyridin-4-yl-1H-benzimidazol-5-yl) amide;
 4-(thiophene-2-sulfonyl)-piperazine-1,3-dicarboxylic acid 3-[(9-ethyl-9H-carbazol-3-yl)-amide] 1-[(2-methoxy-ethyl)-amide];
 4-octyl-1-(thiophene-2-sulfonyl)-piperazine-2-carboxylic acid
 (1-ethyl-2-pyridin-3-yl-1H-benzimidazol-5-yl) amide.

32. A compound according to any one of claims 23 through 31 for use as a medicament.

33. Use of a compound of any one of claim 1 through 31 for preparation of a medicament to treat infertility.

34. A pharmaceutical composition comprising a pharmaceutically acceptable carrier

and one or more compounds of any one of claims 1 through 31.

35. A pharmaceutical composition of claim 34 wherein the compound is packaged together with instructions for use of the compound to treat infertility.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/50640

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D409/14 C07D403/14 C07D403/12 C07D471/04
 //(C07D471/04,221:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, EMBASE, MEDLINE, BIOSIS, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 423 723 B1 (BUCKLER DAVID ET AL) 23 July 2002 (2002-07-23) cited in the application column 4, line 45 -column 6, line 26	1-35
A	BREITENBUCHER J G ET AL: "Generation of a Piperazine-2-carboxamide Library: A Practical Application of the Phenol-Sulfide React and Release Linker" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 39, no. 11, 12 March 1998 (1998-03-12), pages 1295-1298, XP004107918 ISSN: 0040-4039 page 1296 -page 1297; table II -/--	1-35

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

6 February 2004

Date of mailing of the international search report

18/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

vanVoorst tot Voorst, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/50640

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 880 128 A (ROSSMAN RANDALL R ET AL) 9 March 1999 (1999-03-09)	1-33
X	claim 6; examples 21,24,44 ---	34,35
A	WO 98 57960 A (SCHERING CORP ;PHARMACOEPIA INC (US)) 23 December 1998 (1998-12-23)	1-33
X	page 78; examples 202,205,206,220,244 ---	34,35
A	WO 99 26921 A (HAGMANN WILLIAM K ;MAGRIOTIS PLATO A (US); MUMFORD RICHARD A (US);) 3 June 1999 (1999-06-03)	1-33
X	claims 1,17,26 ---	34,35
A	WO 00 39119 A (SCHERING CORP ;PHARMACOEPIA INC (US)) 6 July 2000 (2000-07-06)	1-33
X	claims 1,4 -----	34,35

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 03/50640

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-22
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-22 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/50640

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6423723	B1	23-07-2002	US 6235755 B1 22-05-2001
			US 2002147345 A1 10-10-2002
			AU 5393199 A 28-02-2000
			CA 2339018 A1 17-02-2000
			EP 1380582 A1 14-01-2004
			EP 1102763 A2 30-05-2001
			JP 2002522433 T 23-07-2002
			WO 0008015 A2 17-02-2000
US 5880128	A	09-03-1999	NONE
WO 9857960	A	23-12-1998	AU 753658 B2 24-10-2002
			AU 8058098 A 04-01-1999
			CN 1266433 T 13-09-2000
			EP 0989983 A1 05-04-2000
			HU 0003138 A2 28-06-2001
			JP 2002504144 T 05-02-2002
			NZ 501376 A 28-03-2002
			WO 9857960 A1 23-12-1998
			ZA 9805203 A 05-01-1999
WO 9926921	A	03-06-1999	AU 751950 B2 05-09-2002
			AU 1532799 A 15-06-1999
			CA 2309341 A1 03-06-1999
			EP 1034164 A1 13-09-2000
			JP 2001524465 T 04-12-2001
			WO 9926921 A1 03-06-1999
			US 6645939 B1 11-11-2003
WO 0039119	A	06-07-2000	AU 2158200 A 31-07-2000
			CA 2355744 A1 06-07-2000
			CN 1346355 T 24-04-2002
			EP 1140909 A1 10-10-2001
			JP 2002533455 T 08-10-2002
			WO 0039119 A1 06-07-2000